








56/4/2
MARKING SCHEME
SR. SECONDARY SCHOOL EXAMINATION, 2020
Subject: CHEMISTRY

Q.No.	Expected Answer / Value Points	Distribution of Marks
SECTION - A		
1.	Due to preferential adsorption of common ions from solution / due to electron capture by sol particles during electrodispersion of metal/ due to formulation of electrical double layer.	1
2.	Due to repulsion between the particles of similar charge.	1
3.	Due to preferential adsorption of Γ^- from dispersion medium.	1
4.	By electrophoresis / by mixing two oppositely charged sols / by boiling / by persistent dialysis / by addition of electrolyte.	1
5.	K_2SO_4	1
6.	$NaCN$	1
7.	$C_6H_5CH_2Cl$	1
8.	CH_3-OH	1
9.	Codeine	1
10.	Glycosidic linkage	1
11.	(b)	1
12.	(c)	1
13.	(d)	1
14.	(c)	1
15.	(c)	1
16.	(B)	1
17.	(A)	1
18.	(C)	1
19.	(A)	1
20.	(C)	1
SECTION - B		
21.	$2MnO_2 + 4KOH + O_2 \longrightarrow 2K_2MnO_4 + 2H_2O$ $3MnO_4^{2-} + 4H^+ \longrightarrow 2MnO_4^- + MnO_2 + 2H_2O$ / $MnO_4^{2-} \xrightarrow[\text{oxidation}]{\text{Electrolytic}} MnO_4^- + e^-$	1 1
	OR	
21.	$Cr_2O_7^{2-} + 6Fe^{2+} + 14H^+ \longrightarrow 2Cr^{3+} + 6Fe^{3+} + 7H_2O$ $Cr_2O_7^{2-} + 3Sn^{2+} + 14H^+ \longrightarrow 2Cr^{3+} + 3Sn^{4+} + 7H_2O$	1 1
22.	(i) Reverse osmosis occurs. (ii) Solution shows positive deviation from Raoult's Law.	1 1
23.	(i) Tetracarbonylnickel(0) sp^3 (ii) Hexafluoridocobaltate(III) sp^3d^2	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$

24.	<p>The partial pressure of the gas in vapour phase (p) is directly proportional to the mole fraction of gas(x) in the solution.</p> $p = K_H \cdot x$ $x = \frac{p}{K_H}$ $x = \frac{760}{1.25 \times 10^6}$ $= 6.08 \times 10^{-4}$	<p>1</p> <p>½</p> <p>½</p>
25.	<p>(i) Chemical compounds used for the treatment of stress, and mild or even severe mental diseases. Example : Equanil / meprobamate / luminal (or any other suitable example)</p> <p>(ii) Sodium salts of sulphonated long chain alcohols or hydrocarbons. Example : Sodium Lauryl sulphate / sodium dodecylbenzenesulphonate (or any other suitable example)</p>	<p>½</p> <p>½</p> <p>½</p> <p>½</p>
26.	<p>i)</p> $\begin{array}{c} \text{CHO} \\ \\ (\text{CHOH})_4 \\ \\ \text{CH}_2\text{OH} \end{array} \xrightarrow{\text{Br}_2 \text{ water}} \begin{array}{c} \text{COOH} \\ \\ (\text{CHOH})_4 \\ \\ \text{CH}_2\text{OH} \end{array}$ <p>ii)</p> $\begin{array}{c} \text{CHO} \\ \\ (\text{CHOH})_4 \\ \\ \text{CH}_2\text{OH} \end{array} \xrightarrow[\text{Conc.}]{\text{Oxidation HNO}_3} \begin{array}{c} \text{COOH} \\ \\ (\text{CHOH})_4 \\ \\ \text{COOH} \end{array}$	<p>1</p> <p>1</p>
27.	<p>(a) The metal is converted into its volatile compound which is collected and decomposed to give pure metal.</p> <p>(b) Different components of a mixture are adsorbed to different extent on an adsorbent.</p> <p style="text-align: center;">OR</p> <p>(i) $2\text{Cu}_2\text{S} + 3\text{O}_2 \longrightarrow 2\text{Cu}_2\text{O} + 2\text{SO}_2$ $2\text{Cu}_2\text{O} + \text{Cu}_2\text{S} \longrightarrow 6\text{Cu} + \text{SO}_2$ / $\text{Cu}_2\text{O} + \text{C} \longrightarrow 2\text{Cu} + \text{CO}$</p> <p>(ii) $2[\text{Ag}(\text{CN})_2]^-_{(\text{aq.})} + \text{Zn}_{(\text{s})} \longrightarrow 2\text{Ag}_{(\text{s})} + [\text{Zn}(\text{CN})_4]^{2-}_{(\text{aq.})}$</p>	<p>1</p> <p>1</p> <p>½</p> <p>½</p> <p>1</p>
SECTION - C		
28.	$k = A e^{-E_a/RT}$ $k = (2.5 \times 10^{14} \text{ s}^{-1}) e^{(-25000 \text{ K}/T)}$ $\frac{-E_a}{RT} = \frac{-25000 \text{ K}}{T}$ $\frac{E_a}{R} = 25000 \text{ K}$ $E_a = 25000 \times R$ $= 25000 \times 8.314 \text{ J/mol}$ $= 207850 \text{ J/mol or } 207.85 \text{ kJ/mol}$	<p>½</p> <p>½</p> <p>1</p>

	$t_{\frac{1}{2}} = \frac{0.693}{k}, k = \frac{0.693}{t_{\frac{1}{2}}}$ $k = \frac{0.693}{300 \text{ min}}$ $= 0.00231 \text{ min}^{-1}$	1
29.	<p>(i) Cr^{2+}, because the stable state of chromium is +3 due to t_{2g}^3 configuration.</p> <p>(ii) $\text{Cu}^{+}_{(\text{aq})}$, due to more negative $\Delta_{\text{hyd}}\text{H}^{\circ}$ of $\text{Cu}^{2+}_{(\text{aq})}$ than $\text{Cu}^{+}_{(\text{aq})}$ / It undergoes disproportionation.</p> <p>(iii) Mn^{3+}, because the most stable state of manganese is +2 due to half filled configuration / $3d^5$.</p>	1 1 1
30.	$\Delta T_f = i K_f m$ $\Delta T_f = i \times K_f \times \frac{w_B \times 1000}{M_B \times w_A}$ $2.94 = i \times 4.9 \times \frac{5 \times 1000}{122 \times 35}$ $i = 0.512$ $\alpha = \frac{i - 1}{\frac{1}{n} - 1}$ $\alpha = \frac{0.512 - 1}{\frac{1}{2} - 1}$ $= 0.976$ $= 97.6\%$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 1
31.	<p>(i) Adipic acid and Hexamethylene diamine $\text{HOOC}(\text{CH}_2)_4\text{COOH}$ $n \text{ H}_2\text{N}(\text{CH}_2)_6\text{NH}_2$</p> <p>(ii)</p> <div style="display: flex; justify-content: space-around; align-items: flex-start;"> <div style="text-align: center;"> $\text{HOH}_2\text{C}-\text{CH}_2\text{OH}$ Ethylene glycol (Ethane-1, 2 - diol) </div> <div style="text-align: center;"> $\text{HOOC}-\text{C}_6\text{H}_4-\text{COOH}$ Terephthalic acid (Benzene-1,4 - di carboxylic acid) </div> </div> <p>(iii)</p> <div style="display: flex; justify-content: space-around; align-items: flex-start;"> <div style="text-align: center;"> $\text{CH}_3-\overset{\text{OH}}{\text{CH}}-\text{CH}_2-\text{COOH}$ 3-Hydroxybutanoic acid </div> <div style="text-align: center;"> $\text{CH}_3-\text{CH}_2-\overset{\text{OH}}{\text{CH}}-\text{CH}_2-\text{COOH}$ 3-Hydroxypentanoic acid </div> </div>	$\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$
32.	<p>(i) $\text{C}_6\text{H}_5\text{NH}_2 < (\text{CH}_3)_2\text{NH} < \text{CH}_3\text{NH}_2$</p> <p>(ii) $(\text{CH}_3)_2\text{NH} > \text{CH}_3\text{NH}_2 > (\text{CH}_3)_3\text{N}$</p> <p>(iii) $(\text{C}_2\text{H}_5)_3\text{N} < (\text{C}_2\text{H}_5)_2\text{NH} < \text{C}_2\text{H}_5\text{NH}_2$</p>	1 1 1

33.	<p>i) </p> <p>ii) </p> <p>iii) </p> <p style="text-align: center;">OR</p> <p>a) </p> <p>b) </p>	1 1 1 1
34.	<p>i) $(\text{CH}_3)_3\text{C}-\text{C}(\text{CH}_3)=\text{CHCH}_3$</p> <p>ii) </p> <p>iii) A =  , B = $\text{C}_6\text{H}_5\text{MgBr}$</p> <p style="text-align: center;">OR</p> <p>i) $\text{CH}_3\text{CH}_2\text{CH}=\text{CH}_2 \xrightarrow{\text{HBr / Peroxide}} \text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{-Br}$ $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{-Br} \xrightarrow{\text{NaI / dry acetone}} \text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{-I}$</p> <p>ii) $\text{C}_6\text{H}_6 \xrightarrow[\text{AlCl}_3(\text{anhyd.})]{\text{CH}_3\text{COCl}} \text{C}_6\text{H}_5\text{COCH}_3$</p> <p>iii) $\text{CH}_3\text{CH}_2\text{OH} \xrightarrow{\text{PCl}_5} \text{CH}_3\text{CH}_2\text{Cl} \xrightarrow{\text{KCN}} \text{CH}_3\text{CH}_2\text{CN}$</p>	1 1 $\frac{1}{2} + \frac{1}{2}$ 1 1 1

