

QUESTION PAPER CODE 30/4/2  
**EXPECTED ANSWER/VALUE POINTS**

**SECTION A**

Question numbers 1 to 20 carry 1 mark each.

Question numbers 1 to 10 are multiple choice questions. Choose the correct option.

1. It is being given that the points A(1, 2), B(0, 0) and C(a, b) are collinear. Which of the following relations between a and b is true?

(A)  $a = 2b$                       (B)  $2a = b$                       (C)  $a + b = 0$                       (D)  $a - b = 0$

Sol. (B)  $2a = b$

1

2. In Figure-2, TP and TQ are tangents drawn to the circle with centre at O. If  $\angle POQ = 115^\circ$  then  $\angle PTQ$  is




Fig. 2

(A)  $115^\circ$                       (B)  $57.5^\circ$                       (C)  $55^\circ$                       (D)  $65^\circ$

Sol. (D)  $65^\circ$

1

OR

From an external point Q, the length of the tangent to a circle is 5 cm and the distance of Q from the centre is 8 cm. The radius of the circle is

(A) 39 cm                      (B) 3 cm                      (C)  $\sqrt{39}$  cm                      (D) 7 cm

Sol. (C)  $\sqrt{39}$  cm

1

3. The mean and median of a distribution are 14 and 15 respectively. The value of mode is

(A) 16                      (B) 17                      (C) 18                      (D) 13

Sol. (B) 17

1

4. The equation  $x^2 - 8x + k = 0$  has real and distinct roots if

(A)  $k = 16$                       (B)  $k > 16$                       (C)  $k = 8$                       (D)  $k < 16$

Sol. (D)  $k < 16$

1

5. The first term of an A.P. is 5 and the last term is 45. If the sum of all the terms is 400, the number of terms is

(A) 20 (B) 8 (C) 10 (D) 16

Sol. (D) 16

1

OR

The 9<sup>th</sup> term of the A.P. – 15, –11, –7, ..., 49 is

(A) 32 (B) 0 (C) 17 (D) 13

Sol. (C) 17

1

6. The number of zeroes for a polynomial  $p(x)$  where graph of  $y = p(x)$  is given in Figure-1, is

(A) 3 (B) 4 (C) 0 (D) 5




Fig. 1

Sol. (A) 3

1

7. A bag contains 3 red, 5 black and 7 white balls. A ball is drawn from the bag at random. The probability that the drawn is not black, is

(A)  $\frac{1}{3}$  (B)  $\frac{9}{15}$  (C)  $\frac{5}{10}$  (D)  $\frac{2}{3}$

Sol. (D)  $\frac{2}{3}$

1

8. The value of  $\theta$  for which  $\cos (10^\circ + \theta) = \sin 30^\circ$ , is

(A)  $50^\circ$  (B)  $40^\circ$  (C)  $80^\circ$  (D)  $20^\circ$

Sol. (A)  $50^\circ$

1

9. Point  $P\left(\frac{a}{8}, 4\right)$  is the mid-point of the line segment joining the points  $A(-5, 2)$  and  $B(4, 6)$ . The value of 'a' is

(A) –4 (B) 4 (C) –8 (D) –2

Sol. (A) –4

1

10. The pair of equations,  $x = 0$  and  $x = -4$  has  
 (A) a unique solution (B) no solution  
 (C) infinitely many solutions (D) only solution (0, 0)

Sol. (B) No solution

1

Fill in the blanks in question numbers 11 to 15.

11. The distance between the points (a, b) and (- a, - b) is \_\_\_\_\_.

Sol.  $2\sqrt{a^2 + b^2}$

1

12. If  $\tan A = 1$ , then  $2 \sin A \cos A =$  \_\_\_\_\_.

Sol. 1

1

13.  $\left(\frac{2 + \sqrt{5}}{3}\right)$  is \_\_\_\_\_ number.

Sol. irrational

1

14. A spherical metal ball of radius 8 cm is melted to make 8 smaller identical balls. The radius of each new ball is \_\_\_\_\_ cm.

Sol. 4

1

15. Let  $\Delta ABC \sim \Delta DEF$  and their areas be respectively  $81 \text{ cm}^2$  and  $144 \text{ cm}^2$ . If  $EF = 24 \text{ cm}$ , then length of side  $BC$  is \_\_\_\_\_ cm.

Sol. 18

1

Answer the following question numbers 16 to 20.

16. After how many decimal places will the decimal representation of the rational number  $\frac{229}{2^2 \times 5^7}$  terminate?

Sol. After 7 decimal places

1

17. Given that  $\text{HCF}(120, 160) = 40$ , find  $\text{LCM}(120, 160)$ .

Sol.  $\text{LCM} = \frac{120 \times 160}{40}$

 $\frac{1}{2}$ 

= 480

 $\frac{1}{2}$

18. In Figure-4, AB and CD are common tangents to circle which touch each other at D. If AB = 8 cm, then find the length of CD.




Fig. 4

Sol.  $AC = CD = BC$

 $\frac{1}{2}$ 

$CD = 4 \text{ cm}$

 $\frac{1}{2}$ 

19. Two dice are thrown simultaneously. What is the probability that the sum of the two numbers appearing on the top is 13?

Sol.  $P(E) = 0$

1

20. In Figure-3, a tightly stretched rope of length 20 m is tied from the top of a vertical pole to the ground. Find the height of the pole if the angle made by the rope with the ground is  $30^\circ$ .




Fig. 3

Sol.  $\sin 30^\circ = \frac{AB}{20}$

 $\frac{1}{2}$ 

$AB = 10 \text{ m}$

 $\frac{1}{2}$ 

## SECTION B

Question numbers 21 to 26 carry 2 marks each.

21. Tree Plantation Drive

A group Housing Society has 600 members, who have their houses in the campus and decided to hold a Tree Plantation Drive on the occasion of New Year. Each household was given the choice of planting a sampling of its choice. The number of different types of samplings planted were:

(i) Neem – 125

(ii) Peepal – 165

(iii) Creepers – 50

(iv) Fruit plants – 150

(v) Flowering plants – 110

On the opening ceremony, one of the plants is selected randomly for a prize. After reading the above passage, answer the following questions.

What is the probability that the selected plant is

(i) A fruit plant or a flowering plant?

(ii) Either a Neem plant or a Peepal plant?

**Sol.** Total outcomes = 600

$$(i) P(\text{Fruit plant or a flowering plant}) = \frac{260}{600} \text{ or } \frac{13}{30} \quad 1$$

$$(ii) P(\text{either neem plant or a peepal plant}) = \frac{290}{600} \text{ or } \frac{29}{60} \quad 1$$

**22. Find the mode of the following distribution:**

<b>Classes:</b>	<b>10 – 20</b>	<b>20 – 40</b>	<b>40 – 60</b>	<b>60 – 80</b>	<b>80 – 100</b>
<b>Frequency:</b>	<b>10</b>	<b>8</b>	<b>12</b>	<b>16</b>	<b>4</b>

**Sol.** Model class = 60 – 80  $\frac{1}{2}$

$$\text{Mode} = l + \left( \frac{f_1 - f_0}{2f_1 - f_0 - f_2} \right) \times h = 60 + \left( \frac{16 - 12}{32 - 12 - 4} \right) \times 20 \quad 1$$

$$= 65 \quad \frac{1}{2}$$

**OR**

**From the following distribution, find the median:**

<b>Classes:</b>	<b>500 – 600</b>	<b>600 – 700</b>	<b>700 – 800</b>	<b>800 – 900</b>	<b>900 – 1000</b>
<b>Frequency:</b>	<b>36</b>	<b>32</b>	<b>32</b>	<b>20</b>	<b>30</b>

Median class: 700 – 800  $\frac{1}{2}$

$$\begin{aligned} \text{Median} &= l + \frac{\left(\frac{N}{2} - cf\right)}{f} \times h \\ &= 700 + \frac{75 - 68}{32} \times 100 \\ &= 721.88 \end{aligned}$$

1

 $\frac{1}{2}$ 

23. In Figure-6, a tent is in the shape of a cylinder surmounted by a conical top. The cylindrical part is 2.1 m high and conical part has slant height 2.8 m. Both the parts have same radius 2 m. Find the area of the canvas used to make the tent. (Use  $\pi = \frac{22}{7}$ )




Fig. 6

**Sol.** Area of canvas =  $\pi r(2h + l)$

$$= \frac{22}{7} \times 2(2 \times 2.1 + 2.8)$$

1

$$= 44 \text{ m}^2$$

1

24. Solve for x:

$$8x^2 - 2x - 3 = 0$$

**Sol.**  $8x^2 - 6x + 4x - 3 = 0$

1

$$(4x - 3)(2x + 1) = 0$$

 $\frac{1}{2}$ 

$$x = \frac{3}{4}, x = -\frac{1}{2}$$

 $\frac{1}{2}$

25. The perimeters of two similar triangles are 30 cm and 20 cm respectively. If one side of the first triangle is 9 cm long, find the length of the corresponding side of the second triangle.

Sol. Let the side of other triangle be x cm

$\therefore$  Ratio of perimeters of two similar triangles is equal to ratio of their corresponding sides  $\frac{1}{2}$

$$\therefore \frac{9}{x} = \frac{30}{20} \quad 1$$

$$x = 6 \text{ cm} \quad \frac{1}{2}$$

OR

In Figure-5,  $\Delta PQR$  is right-angled at P. M is a point on QR such that PM is perpendicular to QR. Show that  $PQ^2 = QM \times QR$ .




Fig. 5

$\Delta PQM \sim \Delta RQP$  [By AA similarity] 1

$$\therefore \frac{PQ}{RQ} = \frac{QM}{PQ}$$

$$\Rightarrow PQ^2 = QM \times QR \quad 1$$

26. Evaluate:

$$\frac{\cos 72^\circ}{\sin 18^\circ} + \frac{\sin 11^\circ}{\cos 79^\circ} - \tan 15^\circ \tan 75^\circ$$

Sol.  $\frac{\cos(90^\circ - 18^\circ)}{\sin 18^\circ} + \frac{\sin(90^\circ - 79^\circ)}{\cos 79^\circ} - \tan(90^\circ - 75^\circ) \cdot \tan 75^\circ$   $1 \frac{1}{2}$

$$= \frac{\sin 18^\circ}{\sin 18^\circ} + \frac{\cos 79^\circ}{\cos 79^\circ} - \cot 75^\circ \tan 75^\circ$$

$$= 1 + 1 - 1 = 1 \quad \frac{1}{2}$$

## SECTION C

Question numbers 27 to 34 carry 3 marks each.

27. In Figure-7, two tangents PA and PB are drawn to a circle with centre O from an external point P. Prove that  $\angle APB = 2 \angle OAP$ .




Fig. 7

- Sol.**  $\angle AOB = 180^\circ - \angle APB$  1
- In  $\triangle AOB$ ,  $\angle AOB + \angle OAB + \angle OBA = 180^\circ$  1
- $\Rightarrow 180^\circ - \angle APB + \angle OAB + \angle OBA = 180^\circ$
- $\Rightarrow \angle APB = 2\angle OAB$  1
- 

28. Solve the pair of equations:

$$\frac{2}{x} + \frac{3}{y} = 11, \quad \frac{5}{x} - \frac{4}{y} = -7$$

Hence, find the value of  $5x - 3y$ .

- Sol.**  $\frac{2}{x} + \frac{3}{y} = 11$  ...(i)
- $\frac{5}{x} - \frac{4}{y} = -7$  ...(ii)

On solving equation (i) & (ii)

$$\left. \begin{array}{l} x = 1 \\ \& y = 1/3 \\ \therefore 5x - 3y = 4 \end{array} \right\} \quad \begin{array}{l} 1+1 \\ 1 \end{array}$$

OR

Taxi charges in a city consist of fixed charges and the remainings charges depend upon the distance travelled. For a journey of 10 km, the charge paid is ₹ 75 and for a journey of 15 km, the charge paid is ₹ 110. Find the fixed charge and charges per km. Hence, find the charge of covering a distance of 35 km.

Let fixed charge be ₹ x and charges per km be ₹ y	$\frac{1}{2}$
$x + 10y = 75$ ... (i)	}
$x + 15y = 110$ ... (ii)	
Solve equation (i) & (ii)	1
$x = 5$ & $y = 7$ ]	$\frac{1}{2} + \frac{1}{2}$
$\therefore$ Total charge for 35 km = $x + 35y = ₹ 250$	$\frac{1}{2}$

**29. Construct a triangle with side 5 cm, 6 cm and 7 cm. Now construct another triangle whose side are  $\frac{2}{3}$  times the corresponding sides of the first triangle.**

**Sol.** Correct construction of given triangle 1  
 Correct construction of similar triangle with scale  $2/3$ . 2

**OR**

**Draw a pair of tangents to a circle of radius 3 cm which are inclined to each other at an angle of  $60^\circ$ .**

**Sol.** Correct construction of circle with radius 3 cm. 1  
 Correct construction of two tangents. 2

**30. Prove that  $\sqrt{5}$  is an irrational number.**

**Sol.** Let  $\sqrt{5}$  be a rational number

$$\sqrt{5} = \frac{a}{b} \quad b \neq 0 \quad \text{HCF}(a, b) = 1 \quad \frac{1}{2}$$

$$\Rightarrow 5 = \frac{a^2}{b^2}, \quad a^2 = 5b^2$$

5 divides a 1

Put  $a = 5c$  (for some integer c)

$$\Rightarrow 25c^2 = 5b^2 \Rightarrow b^2 = 5c^2$$

then we get, 5 divides b  $\frac{1}{2}$

Contradiction arises as  $\text{HCF}(a, b) = 1$

$\therefore$  Our assumption is wrong

$\therefore \sqrt{5}$  is irrational number

1

**31. If the sum of the first 6 terms of an A.P. is 36 and that of the first 16 terms is 256, find the sum of the first 11 terms.**

**Sol.** Let a be first term and d be common difference

$$\text{Sum of first 6 terms} = 36 \Rightarrow 2a = 12 - 5d \quad 1$$

$$\text{Sum of first 16 terms} = 256 \Rightarrow 2a = 32 - 15d \quad \frac{1}{2}$$

$$\text{Getting } a = 1, d = 2 \quad 1$$

$$\text{Getting the sum of first 11 terms} = 121 \quad \frac{1}{2}$$

**32. Find the co-ordinates of the points of trisection of the line segment joining the points (3, -1) and (6,8).**

**Sol.**

Case I: If C and D trisect AB



then C divides AB in the ratio 1 : 2  $\frac{1}{2}$

$$\text{Co-ordinates of C: } x = \frac{1 \times 6 + 2 \times 3}{3} = 4 \quad \frac{1}{2}$$

$$\text{and } y = \frac{1 \times 8 + 2(-1)}{3} = 2 \quad \frac{1}{2}$$

$\therefore$  Co-ordinates of C(4, 2)

Case II: Coordinates of D if D divides AB in the ratio 2 : 1  $\frac{1}{2}$

$$\text{Co-ordinates of D: } x' = \frac{2 \times 6 + 1 \times 3}{3} = 5 \quad \frac{1}{2}$$

$$y' = \frac{2 \times 8 + 1 \times (-1)}{3} = 5 \quad \frac{1}{2}$$

Coordinates of D = (5, 5)

OR

Find the area of a quadrilateral ABCD having vertices at A(1, 2), B(1, 0), C(4, 0) and D(4, 4).

$$\text{ar } (\Delta ABC) = \frac{1}{2}[1(0-0) + 1(0-2) + 4(2-0)]$$

$$= 3 \text{ sq. units} \quad 1\frac{1}{2}$$

$$\text{ar } (\Delta ACD) = \frac{1}{2}[1(0-4) + 4(4-2) + 4(2-0)]$$

$$= 6 \text{ sq. units} \quad 1$$

$$\therefore \text{Area of quadrilateral} = 3 + 6 = 9 \text{ sq. units} \quad \frac{1}{2}$$

33. Prove that:

$$\frac{\cos A - \sin A + 1}{\cos A + \sin A - 1} = \operatorname{cosec} A + \cot A$$

Sol. Dividing N<sup>r</sup> & D<sup>r</sup> by sin A in LHS

$$= \frac{\cot A - 1 + \operatorname{cosec} A}{\cot A + 1 - \operatorname{cosec} A} \quad 1$$

$$= \frac{\cot A + \operatorname{cosec} A - (\operatorname{cosec}^2 A - \cot^2 A)}{\cot A + 1 - \operatorname{cosec} A} \quad 1$$

$$= \operatorname{cosec} A + \cot A \quad 1$$

34. In Figure-8, find the area of the shaded region where a circular arc of radius 7 cm has been drawn with vertex O of an equilateral triangle OAB of side 14 cm as centre. (Use  $\pi = \frac{22}{7}$  and  $\sqrt{3} = 1.73$ )




Fig. 8

**Sol.** Area of shaded region =  $\frac{\pi r^2 \theta}{360^\circ} + \frac{\sqrt{3}}{4} a^2$  1

$$= \frac{\pi \times 7^2 \times 300^\circ}{360^\circ} + \frac{\sqrt{3}}{4} \times 14^2$$
1

$$= 213.1 \text{ cm}^2$$
1


---

### SECTION D

**Question numbers 35 to 40 carry 4 marks each.**

- 35. If a line is drawn parallel to one side of a triangle to intersect the other two sides in distinct points, then prove that the other two sides are divided in the same ratio.**

**Sol.** For correct given, To prove, Construction and figure  $4 \times \frac{1}{2} = 2$

For correct proof 2

**OR**

**In a right-angled triangle, prove that the square of the hypotenuse is equal to the sum of the squares of the other two sides.**

**Sol.** For correct given, To prove, construction & figure  $4 \times \frac{1}{2} = 2$

For correct proof 2

---

- 36. A bucket open at the top has top and bottom radii of circular ends as 40 cm and 20 cm respectively. Find the volume of the bucket if its depth is 21 cm. Also find the area of the tin sheet required for making the bucket. (Use  $\pi = \frac{22}{7}$ )**

**Sol.** Volume =  $\frac{\pi h}{3} [R^2 + r^2 + Rr]$

$$= \frac{22}{7} \times \frac{21}{3} [40^2 + 20^2 + 40 \times 20]$$
1

$$= 61600 \text{ cm}^3$$
 $\frac{1}{2}$ 

$$l = \sqrt{h^2 + (R - r)^2} = 29 \text{ cm}$$
1

Area of tin =  $\pi l (R + r) + \pi r^2$

$$= \pi [29 \times 60 + 400]$$
1

$$= 6725.7 \text{ cm}^2$$
 $\frac{1}{2}$ 


---

37. Obtain other zeroes of the polynomial

$$f(x) = 2x^4 + 3x^3 - 5x^2 - 9x - 3$$

if two of its zeroes are  $\sqrt{3}$  and  $-\sqrt{3}$

Sol.  $f(x) = 2x^4 + 3x^3 - 5x^2 - 9x - 3$

$\therefore \sqrt{3}$  and  $-\sqrt{3}$  are zeroes of  $f(x)$

$\therefore (x - \sqrt{3})$  and  $(x + \sqrt{3})$  are factors of  $f(x)$

 $\frac{1}{2}$ 

$\therefore x^2 - 3$  is a factor of  $f(x)$

 $\frac{1}{2}$ 

$$q(x) = \frac{2x^4 + 3x^3 - 5x^2 - 9x - 3}{x^2 - 3}$$

2

$$= 2x^2 + 3x + 1$$

For zeroes  $q(x) = 0$

$$\therefore 2x^2 + 3x + 1 = 0$$

$$(x + 1)(2x + 1) = 0$$

 $\frac{1}{2}$ 

$$x = -1, -1/2$$

$\therefore$  Remaining zeroes are  $-1$  &  $-1/2$

 $\frac{1}{2}$ 

**OR**

Without actually calculating the zeroes, form a quadratic polynomial whose zeroes are reciprocals of the zeroes of the polynomial  $5x^2 + 2x - 3$ .

Let zeroes of given quadratic polynomial be  $\alpha$  and  $\beta$

$$\left. \begin{aligned} \alpha + \beta &= \frac{-2}{5} \\ \alpha\beta &= \frac{-3}{5} \end{aligned} \right\}$$

1

Now,

$$\frac{1}{\alpha} + \frac{1}{\beta} = \frac{\beta + \alpha}{\alpha\beta} = \frac{\frac{-2}{5}}{\frac{-3}{5}} = \frac{2}{3}$$

1

$$\frac{1}{\alpha\beta} = \frac{-5}{3}$$

1

Required Polynomial is

$$x^2 - \frac{2}{3}x - \frac{5}{3}$$

1

or

$$3x^2 - 2x - 5$$

**38. Draw a 'less than ogive for the following distribution. Hence, find median from the graph.**

Marks	Number of Students
0 – 10	2
10 – 20	8
20 – 30	12
30 – 40	10
40 – 50	16
50 – 60	8
60 – 70	3
70 – 80	1

**Sol.** Plotting the points (10, 2), (20, 10), (30, 22)  
(40, 32), (50, 48), (60, 56), (70, 59), (80, 60)

2

Drawing the correct Ogive

 $1\frac{1}{2}$ 

Finding correct Median = 38

 $\frac{1}{2}$ 

**39. In a flight of 600 km, the speed of the aircraft was slowed down due to bad weather. The average speed of the trip was decreased by 200 km/hr and thus the time of flight increased by 30 minutes. Find the average speed of the aircraft originally.**

**Sol.** Let average speed of aircraft be x km/h

$$\frac{600}{x-200} - \frac{600}{x} = \frac{1}{2}$$

2

$$x^2 - 200x - 240000 = 0$$

1

$$(x - 600)(x + 400) = 0$$

$$x = 600 \text{ km/h}$$

1

∴ Original speed = 600 km/h

OR

₹ 9,000 were divided equally among a certain number of persons. Had there been 20 more persons, each would have got ₹ 160 less. Find the original number of persons.

Let original number of persons be  $x$

$$\frac{9000}{x} - \frac{9000}{x+20} = 160 \quad 2$$

$$x^2 + 20x - 1125 = 0 \quad 1$$


$$(x + 45)(x - 25) = 0$$

$$x = 25$$

$$\therefore \text{Number of persons} = 25 \quad 1$$

40. The angle of elevation of an airplane from point A on the ground is  $60^\circ$ . After a flight of 10 seconds, on the same height, the angle of elevation from point A becomes  $30^\circ$ . If the airplane is flying at the speed of 720 km/hr, find the constant height at which the airplane is flying.

Sol.



Correct figure 1

$$\text{Distance travelled in 10 seconds} = 2000 \text{ m} \quad \frac{1}{2}$$

$$\text{Getting } x = \frac{h}{\sqrt{3}} \text{ (In } \triangle ABC) \quad 1$$

$$\text{In } \triangle EDA \tan 30^\circ = \frac{ED}{AD} \Rightarrow \frac{1}{\sqrt{3}} = \frac{h}{x + 2000} \quad 1$$

$$\text{Getting correct value of } h = 1000\sqrt{3} \text{ m.} \quad \frac{1}{2}$$