















MARKING SCHEME
SR. SECONDARY SCHOOL EXAMINATION, 2020
Subject: CHEMISTRY

Q.No.	Expected Answer / Value Points	Distribution of Marks
SECTION - A		
1.	Due to preferential adsorption of common ions from solution / due to electron capture by sol particles during electrodispersion of metal/ due to formulation of electrical double layer.	1
2.	Due to repulsion between the particles of similar charge.	1
3.	Due to preferential adsorption of Γ^- from dispersion medium.	1
4.	By electrophoresis / by mixing two oppositely charged sols / by boiling / by persistent dialysis / by addition of electrolyte.	1
5.	K_2SO_4	1
6.	Distillation / Electrolytic refining	1
7.	$CH_2=CH-CH_2Cl$	1
8.	$(CH_3)_3N$	1
9.	Fibrous Proteins	1
10.	Bithionol / Bithional	1
11.	(c)	1
12.	(a)	1
13.	(b)	1
14.	(c)	1
15.	(b)	1
16.	(D)	1
17.	(C)	1
18.	(D)	1
19.	(A)	1
20.	(B)	1
SECTION – B		
21.	(a) The metal is converted into its volatile compound which is collected and decomposed to give pure metal.	1
	(b) Different components of a mixture are adsorbed to different extent on an adsorbent.	1
OR		
21.	(i) $2Cu_2S + 3O_2 \longrightarrow 2Cu_2O + 2SO_2$ $2Cu_2O + Cu_2S \longrightarrow 6Cu + SO_2 / Cu_2O + C \longrightarrow 2Cu + CO$	$\frac{1}{2}$ $\frac{1}{2}$
	(ii) $2[Ag(CN)_2]^-_{(aq)} + Zn_{(s)} \longrightarrow 2Ag_{(s)} + [Zn(CN)_4]^{2-}_{(aq)}$	1
22.	(i) Reverse osmosis occurs.	1
	(ii) Solution shows positive deviation from Raoult's Law.	1

23.	i) Hexaamminecobalt(III) d^2sp^3 ii) Tetrachloridonickelate (II) sp^3	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$
24.	$2MnO_2 + 4KOH + O_2 \longrightarrow 2K_2MnO_4 + 2H_2O$ $3MnO_4^{2-} + 4H^+ \longrightarrow 2MnO_4^- + MnO_2 + 2H_2O /$ $MnO_4^{2-} \xrightarrow[\text{oxidation}]{\text{Electrolytic}} MnO_4^- + e^-$ OR	1 1
24.	$Cr_2O_7^{2-} + 6Fe^{2+} + 14H^+ \longrightarrow 2Cr^{3+} + 6Fe^{3+} + 7H_2O$ $Cr_2O_7^{2-} + 3Sn^{2+} + 14H^+ \longrightarrow 2Cr^{3+} + 3Sn^{4+} + 7H_2O$	1 1
25.	(i)  (ii) 	1 1
26.	The partial pressure of the gas in vapour phase (p) is directly proportional to the mole fraction of gas(x) in the solution. $p = K_H \cdot x$ $x = \frac{p}{K_H}$ $x = \frac{760}{1.25 \times 10^6}$ $= 6.08 \times 10^{-4}$	1 $\frac{1}{2}$ $\frac{1}{2}$
27.	i) Chemical substances used for the treatment of hyperacidity in the stomach Example: $Al(OH)_3 / Mg(OH)_2 / NaHCO_3$ (or any other suitable example) ii) Chemical substances used to provide sweetness to food with low calories Example: Sucralose / Saccharin / Aspartame (or any other suitable example)	$\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$
SECTION - C		
28.	(i) Cr^{2+} , because the stable state of chromium is +3 due to t_{2g}^3 configuration. (ii) $Cu^+_{(aq)}$, due to more negative $\Delta_{hyd}H^0$ of $Cu^{2+}_{(aq)}$ than $Cu^+_{(aq)}$ / It undergoes disproportionation. (iii) Mn^{3+} , because the most stable state of manganese is +2 due to half filled configuration / $3d^5$.	1 1 1
29.	$\Delta T_f = i K_f m$ $\Delta T_f = i \times K_f \times \frac{w_B \times 1000}{M_B \times w_A}$ $2.94 = i \times 4.9 \times \frac{5 \times 1000}{122 \times 35}$ $i = 0.512$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$

	$\alpha = \frac{i-1}{\frac{1}{n}-1}$ $\alpha = \frac{0.512-1}{\frac{1}{2}-1}$ $= 0.976$ $= 97.6\%$	<p>1/2</p> <p>1</p>
30.	$k = A e^{-E_a/RT}$ $k = (2.5 \times 10^{14} \text{ s}^{-1}) e^{(-25000 \text{ K}/T)}$ $\frac{-E_a}{RT} = \frac{-25000 \text{ K}}{T}$ $\frac{E_a}{R} = 25000 \text{ K}$ $E_a = 25000 \times R$ $= 25000 \times 8.314 \text{ J/mol}$ $= 207850 \text{ J/mol or } 207.85 \text{ kJ/mol}$ $t_{\frac{1}{2}} = \frac{0.693}{k}, k = \frac{0.693}{t_{\frac{1}{2}}}$ $k = \frac{0.693}{300 \text{ min}}$ $= 0.00231 \text{ min}^{-1}$	<p>1/2</p> <p>1/2</p> <p>1</p> <p>1</p>
31.	<p>(a)</p> <p> $\text{CH}_2 = \text{CH} - \text{CH} = \text{CH}_2$ $\text{CH} = \text{CH}_2$   1, 3-Butadiene , Styrene </p> <p>(b)</p> <p> $\text{HOCH}_2 - \text{CH}_2 - \text{OH}$ HOOC COOH   Ethylene glycol , Phthalic acid </p> <p>(c)</p> <p>  , HCHO ; Phenol and formaldehyde </p>	<p>1/2 x 6</p>

32.	<p>i) </p> <p>ii) </p> <p>iii) </p> <p style="text-align: center;">OR</p> <p>a) </p> <p>b) </p>	1 1 1 1 1
33.	<p>i) $(\text{CH}_3)_3\text{C}-\text{C}(\text{CH}_3)=\text{CHCH}_3$</p> <p>ii) </p> <p>iii) A = , B = $\text{C}_6\text{H}_5\text{MgBr}$</p> <p style="text-align: center;">OR</p> <p>i) $\text{CH}_3\text{CH}_2\text{CH}=\text{CH}_2 \xrightarrow{\text{HBr / Peroxide}} \text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{-Br}$ $\downarrow \text{NaI / dry acetone}$ $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{-I}$</p> <p>ii) $\text{C}_6\text{H}_6 \xrightarrow[\text{AlCl}_3(\text{anhyd.})]{\text{CH}_3\text{COCl}} \text{C}_6\text{H}_5\text{COCH}_3$</p> <p>iii) $\text{CH}_3\text{CH}_2\text{OH} \xrightarrow{\text{PCl}_5} \text{CH}_3\text{CH}_2\text{Cl} \xrightarrow{\text{KCN}} \text{CH}_3\text{CH}_2\text{CN}$</p>	1 1 $\frac{1}{2} + \frac{1}{2}$ 1 1 1
34.	<p>(i) $\text{C}_6\text{H}_5\text{NH}_2 < (\text{CH}_3)_2\text{NH} < \text{CH}_3\text{NH}_2$</p> <p>(ii) $(\text{CH}_3)_2\text{NH} > \text{CH}_3\text{NH}_2 > (\text{CH}_3)_3\text{N}$</p> <p>(iii) $(\text{C}_2\text{H}_5)_3\text{N} < (\text{C}_2\text{H}_5)_2\text{NH} < \text{C}_2\text{H}_5\text{NH}_2$</p>	1 1 1

