

Marking Scheme
Strictly Confidential
(For Internal and Restricted use only)
Senior Secondary School Certificate Examination,2024
SUBJECT NAME BIOLOGY (Q.P. CODE 57/4/3)


General Instructions: -


1	You are aware that evaluation is the most important process in the actual and correct assessment of the candidates. A small mistake in evaluation may lead to serious problems which may affect the future of the candidates, education system and teaching profession. To avoid mistakes, it is requested that before starting evaluation, you must read and understand the spot evaluation guidelines carefully.
2	“Evaluation policy is a confidential policy as it is related to the confidentiality of the examinations conducted, Evaluation done and several other aspects. Its’ leakage to public in any manner could lead to derailment of the examination system and affect the life and future of millions of candidates. Sharing this policy/document to anyone, publishing in any magazine and printing in News Paper/Website etc may invite action under various rules of the Board and IPC.”
3	Evaluation is to be done as per instructions provided in the Marking Scheme. It should not be done according to one’s own interpretation or any other consideration. Marking Scheme should be strictly adhered to and religiously followed. However, while evaluating, answers which are based on latest information or knowledge and/or are innovative, they may be assessed for their correctness otherwise and due marks be awarded to them. In class-XII, while evaluating two competency-based questions, please try to understand given answer and even if reply is not from marking scheme but correct competency is enumerated by the candidate, due marks should be awarded.
4	The Marking scheme carries only suggested value points for the answers These are in the nature of Guidelines only and do not constitute the complete answer. The students can have their own expression and if the expression is correct, the due marks should be awarded accordingly.
5	The Head-Examiner must go through the first five answer books evaluated by each evaluator on the first day, to ensure that evaluation has been carried out as per the instructions given in the Marking Scheme. If there is any variation, the same should be zero after deliberation and discussion. The remaining answer books meant for evaluation shall be given only after ensuring that there is no significant variation in the marking of individual evaluators.
6	Evaluators will mark(√) wherever answer is correct. For wrong answer CROSS ‘X’ be marked. Evaluators will not put right (✓) while evaluating which gives an impression that answer is correct and no marks are awarded. This is most common mistake which evaluators are committing.
7	If a question has parts, please award marks on the right-hand side for each part. Marks awarded for different parts of the question should then be totaled up and written in the left-hand margin and encircled. This may be followed strictly.

8	If a question does not have any parts, marks must be awarded in the left-hand margin and encircled. This may also be followed strictly.
9	If a student has attempted an extra question, answer of the question deserving more marks should be retained and the other answer scored out with a note “ Extra Question ”.
10	No marks to be deducted for the cumulative effect of an error. It should be penalized only once.
11	A full scale of marks 70 has to be used. Please do not hesitate to award full marks if the answer deserves it.
12	Every examiner has to necessarily do evaluation work for full working hours i.e., 8 hours every day and evaluate 20 answer books per day in main subjects and 25 answer books per day in other subjects (Details are given in Spot Guidelines). This is in view of the reduced syllabus and number of questions in question paper.
13	<p>Ensure that you do not make the following common types of errors committed by the Examiner in the past:-</p> <ul style="list-style-type: none"> ● Leaving answer or part thereof unassessed in an answer book. ● Giving more marks for an answer than assigned to it. ● Wrong totaling of marks awarded on an answer. ● Wrong transfer of marks from the inside pages of the answer book to the title page. ● Wrong question wise totaling on the title page. ● Wrong totaling of marks of the two columns on the title page. ● Wrong grand total. ● Marks in words and figures not tallying/not same. ● Wrong transfer of marks from the answer book to online award list. ● Answers marked as correct, but marks not awarded. (Ensure that the right tick mark is correctly and clearly indicated. It should merely be a line. Same is with the X for incorrect answer.) ● Half or a part of answer marked correct and the rest as wrong, but no marks awarded.
14	While evaluating the answer books if the answer is found to be totally incorrect, it should be marked as cross (X) and awarded zero (0) Marks.
15	Any unassessed portion, non-carrying over of marks to the title page, or totaling error detected by the candidate shall damage the prestige of all the personnel engaged in the evaluation work as also of the Board. Hence, in order to uphold the prestige of all concerned, it is again reiterated that the instructions be followed meticulously and judiciously.
16	The Examiners should acquaint themselves with the guidelines given in the “ Guidelines for Spot Evaluation ” before starting the actual evaluation.
17	Every Examiner shall also ensure that all the answers are evaluated, marks carried over to the title page, correctly totaled and written in figures and words.
18	The candidates are entitled to obtain photocopy of the Answer Book on request on payment of the prescribed processing fee. All Examiners/Additional Head Examiners/Head Examiners are once again reminded that they must ensure that evaluation is carried out strictly as per value points for each answer as given in the Marking Scheme.

MARKING SCHEME
Senior Secondary School Examination, 2024
BIOLOGY (Subject Code-044)
[Paper Code: 57/4/3]


SECTION-A			
1.	(A)/ antipodal, zygote and endosperm	1	1
2.	(A)/ A – connective, B – Endothecium, C – Pollen grain	1	1
3.	(C)/ Autosomal recessive	1	1
4.	(D)/ P – Transcription, Q – mRNA, R – Translation	1	1
5.	(B)/ Evening primrose	1	1
6.	(C)/ A - (ii), B – (iv), C – (iii), D – (v)	1	1
7.	(C)/ A - III, B - I, C - IV, D - II	1	1
8.	(C)/ Ovum - A, Morula - E, Blastocyst - G	1	1
9.	(C)/ BAC and YAC	1	1
10.	(A)/ Aneuploidy	1	1
11.	(B)/ Antigen- Antibody interaction	1	1
12.	(D)/ Restriction Enzymes	1	1
13.	(C)/ (A) is true, but (R) is false.	1	1
14.	(A)/ Both Assertion (A) and Reason (R) are true and Reason (R) is correct explanation of Assertion (A).	1	1
15.	(B)/ Both Assertion (A) and Reason (R) are true and Reason (R) is not correct explanation of Assertion (A).	1	1
16.	(A)/ Both Assertion (A) and Reason (R) are true and Reason (R) is the correct explanation of Assertion (A).	1	1
SECTION - B			
17.	(a) Rosie is the first transgenic cow (b) The milk contained human protein alpha-lactoalbumin (2.4 g per liter) which is nutritionally a more balanced product than natural cow-milk.	1 1	2
18.	<ul style="list-style-type: none"> • X- In woman X thickness of uterine wall increases after mid of menstrual cycle; Reason- due to fertilization of egg/pregnancy/conceived • Y- In woman Y thickness of uterine wall decreases after mid of menstrual cycle; Reason- egg has not been fertilized/leading to the breakdown of lining of the uterus/menstrual flow/ bleeding. 	1/2+1/2 1/2+1/2	2
19.	(a) Sharing of injection needles or surgical instruments etc. with an infected person, Transfusion of blood contaminated with HIV/Hepatitis B virus, An infected mother to the foetus through placenta [Any Two]	1/2+1/2	
	(b) For proper treatment and to prevent spread of the disease.	1	2
20.	(a) Biodiversity hotspots are the regions with very high level of species richness, and high degree of endemism	1/2+1/2	
	(b) Western Ghats, The Himalayas, Indo – Burma region [Any two]	1/2+1/2	2

21.	(a)														
<table border="1"> <thead> <tr> <th data-bbox="256 298 483 373"></th> <th data-bbox="483 298 852 373">Grazing Food Chain</th> <th data-bbox="852 298 1234 373">Detritus Food Chain</th> </tr> </thead> <tbody> <tr> <td data-bbox="256 373 483 489">1.</td> <td data-bbox="483 373 852 489">It starts with producers.</td> <td data-bbox="852 373 1234 489">Starts from dead organic matter/ detritus</td> </tr> <tr> <td data-bbox="256 489 483 604">2.</td> <td data-bbox="483 489 852 604">2nd trophic level is occupied by herbivores.</td> <td data-bbox="852 489 1234 604">2nd trophic level is occupied by detritivores.</td> </tr> <tr> <td data-bbox="256 604 483 835">3.</td> <td data-bbox="483 604 852 835">In aquatic ecosystem a major fraction of energy flows through this type of food chain</td> <td data-bbox="852 604 1234 835">In terrestrial ecosystem much larger fraction of energy flow through such type of food chain.</td> </tr> </tbody> </table> <p data-bbox="776 835 1214 877" style="text-align: center;">(Any two corresponding points)</p> <p data-bbox="722 945 776 982" style="text-align: center;">OR</p> <p data-bbox="256 982 300 1020">(b)</p> <ul data-bbox="305 1020 1234 1239" style="list-style-type: none"> • Brood parasitism is a method of parasitism in which the parasitic bird lays its eggs in the nest of the host bird and let the host bird to incubate them. • Example – The cuckoo bird lays its eggs in the nest of crow, egg of cuckoo bird resemble crows egg in size and colour to reduce the chance of host bird detecting and ejecting them from nest 			Grazing Food Chain	Detritus Food Chain	1.	It starts with producers.	Starts from dead organic matter/ detritus	2.	2 nd trophic level is occupied by herbivores.	2 nd trophic level is occupied by detritivores.	3.	In aquatic ecosystem a major fraction of energy flows through this type of food chain	In terrestrial ecosystem much larger fraction of energy flow through such type of food chain.	1+1	
	Grazing Food Chain	Detritus Food Chain													
1.	It starts with producers.	Starts from dead organic matter/ detritus													
2.	2 nd trophic level is occupied by herbivores.	2 nd trophic level is occupied by detritivores.													
3.	In aquatic ecosystem a major fraction of energy flows through this type of food chain	In terrestrial ecosystem much larger fraction of energy flow through such type of food chain.													
SECTION - C															
22.	 <p data-bbox="344 1774 641 1806" style="color: red;">[1/2 mark for each labeling]</p>	1/2 x6													
			3												

27.	<p>Phenotype of parents → Purple flower × White flower</p> <p>Genotype → PP [1/2 mark]</p> <p>Gametes → (P) (p)</p> <p>F₁ generation → Pp [1/2 mark] (50) Purple flower</p> <p>Selfing of F₁ generation → Pp × Pp</p> <p>Gametes → (P) (p) (P) (p)</p> <p>F₂ generation →</p> <table border="1" data-bbox="617 525 893 693"> <tr> <td>♀ \ ♂</td> <td>P</td> <td>p</td> <td></td> </tr> <tr> <td>P</td> <td>PP Purple</td> <td>Pp Purple</td> <td>[1/2 mark]</td> </tr> <tr> <td>p</td> <td>Pp Purple</td> <td>pp White</td> <td></td> </tr> </table> <p>Purple : White 482 : 162 3 : 1 [1/2 mark]</p> <p>- It is due to complete dominance</p>	♀ \ ♂	P	p		P	PP Purple	Pp Purple	[1/2 mark]	p	Pp Purple	pp White		1/2 x4	1	3
♀ \ ♂	P	p														
P	PP Purple	Pp Purple	[1/2 mark]													
p	Pp Purple	pp White														
28.	 <p>[Vegetative cell, generative cell, exine, intine, germ pore]</p> <p>(Any four labelings to be considered)</p> <p>Vegetative cell- contain abundant food reserve for pollen germination Generative cell- Divide to produce male gametes Exine – Protective layer Intine- Involved in the formation of pollen tube. Germ pore- Pollen tube emerges from this point.</p> <p>[Marks to be given to two correct labelings with their respective correct function only]</p>	1/2 x4	1/2 x2	3												
SECTION - D																
29.	<p>(a)</p> <ul style="list-style-type: none"> • hnRNA/ heterogeneous nuclear RNA • RNA polymerase II <p>(b) hnRNA undergo capping at 5' end (methyl guanosine triphosphate /mGppp), and tailing at 3' end (with poly A tail or adenylate residue), further splicing is carried out, where non coding sequences or introns are removed and coding sequence or exons are joined together/ diagrammatic</p>	1/2 1/2	1/2x4													

	<p>representation with given markers can also be considered.</p> <p>(c)</p> <ul style="list-style-type: none"> In prokaryotes - 1 In eukaryotes-3 <p style="text-align: center;">OR</p> <p>(c) In prokaryotes the transcription takes place in the cytoplasm/cytosol whereas in Eukaryote transcription occurs in the Nucleus,</p>	<p>1/2</p> <p>1/2</p> <p>1</p>	<p>4</p>
30.	<p>(a)</p> <ul style="list-style-type: none"> A- exponential growth curve/ J-shaped curve B- Logistic growth curve/ S- shaped curve/ sigmoid curve <p>(b)</p> <ul style="list-style-type: none"> It represents carrying capacity (K) Maximum possible number of individuals beyond which no growth of population is observed. <p style="text-align: center;">OR</p> <p>(b)</p> <p>Growth curve B is formed when resources (food and space) in nature are limited have environmental checks while growth curve A is formed when resources are unlimited with no environmental checks.</p> <p>(c)</p> <p>(i)</p> <ul style="list-style-type: none"> 'B'/ Logistic growth curve As resources are never infinite in nature. <p>(ii)</p> <ul style="list-style-type: none"> J-shaped curve/ exponential growth It is a continuous growing population. 	<p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p>	<p>4</p>
SECTION - E			
31.	<p>(a)</p> <p>(i)</p> <ul style="list-style-type: none"> X to \bar{X} is 5' \longrightarrow 3' No more amino acids will be added as the last codon UAA is a stop codon <p>(ii)</p> <ul style="list-style-type: none"> AUG Anticodon - UAC Methionine <p>(iii)</p> <ul style="list-style-type: none"> The amino acids are activated in the presence of ATP , and linked to their cognate tRNA or the adapter molecule, 	<p>1/2 x3</p> <p>1/2 x3</p> <p>1/2+1/2</p>	

	<ul style="list-style-type: none"> Amino acids are activated so peptide bonds can be formed using this energy. <p style="text-align: center;">OR</p> <p>(b)</p> <p>(i) the RBC in such patients takes up a sickle shape instead of biconcave.</p> <p>(ii) The defect is caused by the substitution of Glutamic acid (Glu) by Valine (Val) ,at the sixth position of the beta globin chain of the haemoglobin molecule /The substitution of amino acid in the globin protein results due to the single base substitution at the sixth codon of the beta globin gene, from GAG to GUG.</p> <p>(iii)</p> <p>Phenotype: ♀ carrier X carrier ♂ Genotype: Hb^A Hb^S crossed with Hb^A Hb^S [1/2 mark]</p> <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td style="text-align: center;">gametes</td> <td style="text-align: center;">Hb^A ♀</td> <td style="text-align: center;">Hb^S ♂</td> <td></td> </tr> <tr> <td style="text-align: center;">Hb^A</td> <td style="text-align: center;">Hb^AHb^A</td> <td style="text-align: center;">Hb^AHb^S</td> <td rowspan="2" style="vertical-align: middle;">[1/2 mark]</td> </tr> <tr> <td style="text-align: center;">♀ Hb^S</td> <td style="text-align: center;">Hb^AHb^S</td> <td style="text-align: center;">Hb^SHb^S</td> </tr> </table> <p>Genotypes: Hb^AHb 2 Hb^AHb^S Hb^SHb^S [1/2mark] Phenotypes: Unaffected Carrier Sickle cell</p> <p>It is an autosomal recessive disorder [1/2 mark]</p>	gametes	Hb ^A ♀	Hb ^S ♂		Hb ^A	Hb ^A Hb ^A	Hb ^A Hb ^S	[1/2 mark]	♀ Hb ^S	Hb ^A Hb ^S	Hb ^S Hb ^S	<p>1</p> <p>1</p> <p>1+1</p> <p>1/2 x4</p>	<p>5</p>
gametes	Hb ^A ♀	Hb ^S ♂												
Hb ^A	Hb ^A Hb ^A	Hb ^A Hb ^S	[1/2 mark]											
♀ Hb ^S	Hb ^A Hb ^S	Hb ^S Hb ^S												
<p>32.</p>	<p>(a) HIV enters human body and enter macrophage cells , where viral RNA is introduced in the cell, Viral DNA is produced by reverse transcription, Viral DNA incorporated in host genome, New viral RNA is produced by the infected cell, New viruses are produced, which enter T-helper cells replicate and produces its progeny, Progeny viruses attack other T helper cells, this is repeated leading to progressive decline in the number T helper cells, person become immunodeficient and HIV AIDS develop./</p>	<p>1/2x10</p>												



New progeny viruses infect other helper T-cells [1/2 mark] → Progressive decline in helper T cells [1/2 mark] → Patient become immuno deficient [1/2 mark]

1/2x10

(b)

(i)


- Chills, and high fever occurs in a cyclic order/every 2-3 days,
- fever is due to the toxic haemozoin,
- release at the time of RBC rupture.

1/2+1/2

1/2

1

(ii)



1/2x5

5

33.	<p>(a)</p> <p>(i) To combine desirable characters from two different varieties , so as to obtain improved/ superior variety</p> <p>(ii) If the female parent plant bears bisexual flowers, it is necessary to remove the anthers from the flower before its dehiscence. which is called emasculation , such flowers are then bagged , (with butter paper bags of suitable size) to prevent contamination of its stigma. which is called bagging , When the stigma of bagged flower attains receptivity, mature pollen collected from anthers of the male parent are dusted on the stigma , and flower is rebagged and fruit is allowed to develop.</p> <p style="text-align: center;">OR</p> <p>(b)</p> <p>(i) After implantation, finger-like projections appear on the trophoblast called chorionic villi , which are surrounded by uterine tissue and maternal blood , The chorionic villi and uterine tissue become interdigitated with each other , and jointly form a structural and functional unit between developing embryo and maternal body called placenta.</p> <p>(ii) Placenta facilitates supply of O₂ and nutrition to the embryo , and removal of CO₂ and waste from foetus , Placenta also acts as endocrine tissue and produces several hormones hCG hPL estrogen progesterogen essential for maintaining pregnancy.</p>	<p>1+1</p> <p>½ x6</p> <p>½ x4</p> <p>1x3</p>	<p>5</p>
-----	--	---	----------