

QUESTION PAPER CODE 430/3/2  
**EXPECTED ANSWER/VALUE POINTS**

**SECTION A**

Question numbers 1 to 10 are multiple choice questions of 1 mark each.

Select the correct choice.

1. The simplest form of  $\frac{1095}{1168}$  is

(a)  $\frac{17}{26}$                       (b)  $\frac{25}{26}$                       (c)  $\frac{13}{16}$                       (d)  $\frac{15}{16}$

Sol. (d)  $\frac{15}{16}$

1

2. One card is drawn at random from a well – shuffled deck of 52 cards. What is the probability of getting a Jack?

(a)  $\frac{3}{26}$                       (b)  $\frac{1}{52}$                       (c)  $\frac{1}{13}$                       (d)  $\frac{3}{52}$

Sol. (c)  $\frac{1}{13}$

1

3. Which of the following rational numbers is expressible as a terminating decimal?

(a)  $\frac{124}{165}$                       (b)  $\frac{131}{30}$                       (c)  $\frac{2027}{625}$                       (d)  $\frac{1625}{462}$

Sol. (c)  $\frac{2027}{625}$

1

4. If one zero of the quadratic polynomial,  $(k - 1)x^2 + kx + 1$  is  $-4$  then the value of  $k$  is

(a)  $-\frac{5}{4}$                       (b)  $\frac{5}{4}$                       (c)  $-\frac{4}{3}$                       (d)  $\frac{4}{3}$

Sol. (b)  $\frac{5}{4}$

1

5. If  $P(-1, 1)$  is the midpoint of the line segment joining  $A(-3, b)$  and  $B(1, b + 4)$ , then  $b$  is equal to

(a) 1                      (b)  $-1$                       (c) 2                      (d) 0

Sol. (b)  $-1$

1

6. Consider the following distribution:

Classes:	0 – 5	5 – 10	10 – 15	15 – 20	20 – 25
Frequency:	10	15	12	20	9

The sum of lower limits of the median class and the modal class is

- (a) 15                                      (b) 25                                      (c) 30                                      (d) 35

Sol. (b) 25

1

7. What is the largest number that divides 245 and 1029, leaving remainder 5 in each?

- (a) 15                                      (b) 16                                      (c) 9                                      (d) 5

Sol. (b) 16

1

8. The distance between the points A(2, –3) and B(2, 2) is

- (a) 2 units                                      (b) 3 units                                      (c) 4 units                                      (d) 5 units

Sol. (d) 5 units

1

9. The product of the two zeroes of the polynomial  $3x^2 - 7x - 27$  is:

- (a) 27                                      (b) 9                                      (c) –9                                      (d)  $\frac{7}{3}$

Sol. (c) –9

1

10. If the tangents PA and PB from an external point P to a circle with centre O are inclined to each other at an angle of  $80^\circ$ , then  $\angle POA$  equals:

- (a)  $50^\circ$                                       (b)  $60^\circ$                                       (c)  $70^\circ$                                       (d)  $80^\circ$

Sol. (a)  $50^\circ$

1

In Question numbers 11 to 15, fill in the blanks:

11. The value of k for which system of equations  $x + 2y = 3$  and  $5x + ky = 7$  has no solution is \_\_\_\_\_.

Sol.  $k = 10$

1

12. The value of  $(\tan 27^\circ - \cot 63^\circ)$  is \_\_\_\_\_.

Sol. 0

1

13. If ratio of the corresponding sides of two similar triangles is 2:3, then ratio of their perimeters is \_\_\_\_\_.

Sol. 2 : 3

1

14. Distance between (a, –b) and (a, b) is \_\_\_\_\_.

Sol. 2b units

1

15. The value of  $(\sin 20^\circ - \cos 70^\circ)$  is \_\_\_\_\_.

Sol. 0

1

Answer the following questions, Question numbers 16 to 20.

16. The perimeter of a sector of a circle of radius 14 cm is 68 cm. Find the area of the sector.

Sol.  $l = 68 - 28 = 40$  cm  $\frac{1}{2}$

$A = 280$  cm<sup>2</sup>  $\frac{1}{2}$

OR

The circumference of a circle is 39.6 cm. Find its area.

Sol.  $r = \frac{39.6}{2\pi}$   $\frac{1}{2}$

$A = \frac{392.04}{\pi}$  or 124.74 cm<sup>2</sup>  $\frac{1}{2}$

17. If  $3y - 1$ ,  $3y + 5$  and  $5y + 1$  are three consecutive terms of an A.P., then find the value of  $y$ .

Sol.  $2(3y + 5) = 3y - 1 + 5y + 1$   $\frac{1}{2}$

$y = 5$   $\frac{1}{2}$

18. If  $\sec \theta = \frac{25}{7}$ , then find the value of  $\cot \theta$ .

Sol.  $\tan \theta = \frac{24}{7} \Rightarrow \cot \theta = \frac{7}{24}$   $\frac{1}{2} + \frac{1}{2}$

OR

If  $3 \tan \theta = 4$ , then find the value of  $\left( \frac{3 \sin \theta + 2 \cos \theta}{3 \sin \theta - 2 \cos \theta} \right)$

Sol. Given expression =  $\frac{3 \times \frac{4}{3} + 2}{3 \times \frac{4}{3} - 2} = 3$   $\frac{1}{2} + \frac{1}{2}$

19. A bag contains 5 red, 4 blue and 3 green balls. A ball is drawn at random from the bag. Find the probability of getting a ball not of blue colour.

Sol. Total No. of balls = 12  $\frac{1}{2}$

$P(\text{Not a blue ball}) = \frac{8}{12}$  or  $\frac{2}{3}$   $\frac{1}{2}$

20. In Fig. 1,  $DE \parallel BC$ ,  $AD = 2.4$  cm,  $AE = 3.2$  cm and  $CE = 4.8$  cm. Find  $BD$




Fig. 1

$$\begin{aligned} \text{Sol. } \frac{2.4}{BD} &= \frac{3.2}{4.8} && \frac{1}{2} \\ \Rightarrow BD &= 3.6 \text{ cm} && \frac{1}{2} \end{aligned}$$

## SECTION B

Question numbers 21 to 26 carry 2 marks each.

21. Prove that:  $\sqrt{\sec^2 \theta + \operatorname{cosec}^2 \theta} = \tan \theta + \cot \theta$

$$\begin{aligned} \text{Sol. LHS} &= \sqrt{\sec^2 \theta + \operatorname{cosec}^2 \theta} = \sqrt{1 + \tan^2 \theta + 1 + \cot^2 \theta} && 1 \\ &= \sqrt{\tan^2 \theta + \cot^2 \theta + 2} \\ &= \sqrt{\tan^2 \theta + \cot^2 \theta + 2 \tan \theta \cot \theta} \\ &= \sqrt{(\tan \theta + \cot \theta)^2} && \frac{1}{2} \\ &= \tan \theta + \cot \theta = \text{RHS} && \frac{1}{2} \end{aligned}$$

OR

Prove that:  $\frac{\sin \theta}{1 - \cos \theta} = (\operatorname{cosec} \theta + \cot \theta)$

$$\begin{aligned} \text{Sol. LHS} &= \frac{\sin \theta}{1 - \cos \theta} \times \frac{1 + \cos \theta}{1 + \cos \theta} && 1 \\ &= \frac{\sin \theta (1 + \cos \theta)}{1 - \cos^2 \theta} && \frac{1}{2} \\ &= \frac{\sin \theta (1 + \cos \theta)}{\sin^2 \theta} = \frac{1}{\sin \theta} + \frac{\cos \theta}{\sin \theta} \\ &= \operatorname{cosec} \theta + \cot \theta = \text{RHS} && \frac{1}{2} \end{aligned}$$

22. Find the values of  $p$  for which the quadratic equation  $x^2 - 2px + 1 = 0$  has no real roots.

Sol. For no real roots

$$D < 0$$

$$(-2p)^2 - 4 \times 1 \times 1 < 0 \quad 1$$

$$p^2 - 1 < 0 \quad \frac{1}{2}$$

$$-1 < p < 1 \quad \frac{1}{2}$$


---

23. Two dice are thrown at the same time. Find the probability of getting different numbers on the two dice.

Sol. Total number of outcomes = 36  $\frac{1}{2}$

Favourable numbers of outcomes = 30  $\frac{1}{2}$

$$\text{Probability} = \frac{30}{36} \text{ or } \frac{5}{6} \quad 1$$

(Both no. are different)

**OR**

Two dice are thrown at the same time. Find the probability that the sum of the two numbers appearing on the top of the dice is more than 9.

Sol. Favourable outcomes (5, 5), (4, 6), (6, 4), (6, 5), (5, 6), (6, 6)

Total number of outcomes = 36  $\frac{1}{2}$

Number of favourable outcomes = 6  $\frac{1}{2}$

$$\text{Required probability} = \frac{6}{36} \text{ or } \frac{1}{6} \quad 1$$


---

24. A bag contains 5 red, 8 white and 7 black balls. A ball is drawn at random from the bag. Find the probability that the drawn ball is

(i) red or white

(ii) not a white ball

Sol. Total no. of balls = 20


$$(i) \text{ P(ball is red or white)} = \frac{13}{20} \quad 1$$

$$(ii) \text{ P(Not a white ball)} = \frac{12}{20} \text{ or } \frac{3}{5} \quad 1$$


---

25. The length of a tangent from a point A at a distance of 5 cm from the centre of the circle is 4 cm. Find the diameter of the circle.

Sol.



$$OA = 5 \text{ cm}$$

$$AB = 4 \text{ cm}$$

$$OB^2 = OA^2 - AB^2$$

$$= 25 - 16$$

$$= 9$$

$$OB = 3 \text{ cm}$$

$$\text{diameter} = 6 \text{ cm}$$

 $\frac{1}{2}$ 

1

 $\frac{1}{2}$ 

26. Find the area of a circle whose circumference is 44 cm.

Sol.  $2\pi r = 44$

$$r = 7 \text{ cm}$$

$$\text{Area of circle} = \frac{22}{7} \times 7 \times 7 = 154 \text{ cm}^2$$

1

1

### SECTION C

Question numbers 27 to 34 carry 3 marks each.

27. In Fig. 3, arrangement of desks in a classroom is shown. Ashima, Bharti and Asha are seated at A, B and C respectively. Answer the following:

(i) Find whether the girls are sitting in a line.

(ii) If A, B and C are collinear, find the ratio in which point B divides the line segment joining A and C.




Fig. 3

**Sol.** Coordinates of A(3, 1)

B(6, 4)

C(8, 6)

$$(i) \text{ Area of } (\Delta ABC) = \frac{1}{2}[3(4-6) + 6(6-1) + 8(1-4)]$$

$$= 0$$

Yes they are sitting in same line

(ii) Let AB:BC = k : 1

$$6 = \frac{8k+3}{k+1}$$

$$k = \frac{3}{2} \text{ or Ratio} = 3:2$$

**28. A number consists of two digits whose sum is 10. If 18 is subtracted from the number, its digit are reversed. Find the number.**

**Sol.** Let two digit number =  $10x + y$

$$x + y = 10 \quad \dots(i)$$

$$10x + y - 18 = 10y + x$$

$$\Rightarrow x - y = 2 \quad \dots(ii)$$

On solving (i) & (ii)  $x = 6, y = 4$

$\therefore$  Required number = 64

**29. If  $\sqrt{2}$  is given as an irrational number, then prove that  $(7 - 2\sqrt{2})$  is an irrational number.**

**Sol.** Let  $7 - 2\sqrt{2} = m$ , where m is a rational number

$$\sqrt{2} = \frac{7-m}{2}$$

Irrational = Rational

$\Rightarrow$  LHS  $\neq$  RHS

It means out assumption is wrong.

Hence,  $7 - 2\sqrt{2}$  is irrational

OR

Find HCF of 44, 96 and 404 by prime factorization method. Hence find their LCM.

<b>Sol.</b>	$44 = 2^2 \times 11$	$\left. \begin{array}{l} \\ \\ \\ \end{array} \right\} \frac{1}{2}$
	$96 = 2^5 \times 3$	
	$404 = 2^2 \times 101$	
	HCF = $2^2 = 4$	$\frac{1}{2}$
	LCM = $2^5 \times 11 \times 3 \times 101$	
	= 106656	1

---

30. If 1 and -2 are the zeroes of the polynomial  $(x^3 - 4x^2 - 7x + 10)$ , find its third zero.

<b>Sol.</b>	The two factors of polynomials are $(x - 1)$ , $(x + 2)$	$\frac{1}{2}$
	$(x - 1)(x + 2) = x^2 + x - 2$	$\frac{1}{2}$
	$\frac{x^3 - 4x^2 - 7x + 10}{x^2 + x - 2} = (x - 5)$	$\frac{1}{2}$
	Third zero = 5	$\frac{1}{2}$

---

31. Draw a circle of radius 3 cm. From a point 7 cm away from its centre, construct a pair of tangents to the circle.

<b>Sol.</b>	Drawing a circle of radius 3 cm, marking Centre O and taking a point P such that OP = 7 cm	$\left. \begin{array}{l} \\ \\ \\ \end{array} \right\} 1$
	Constructing two tangents	2

OR

Draw a line segment of 8 cm and divide it in the ratio 3 : 4.

<b>Sol.</b>	Drawing a line segment of 8 cm	1
	Dividing it in the ratio 3 : 4	2

---

32. Prove that  $\frac{\cos \theta}{(1 - \tan \theta)} + \frac{\sin \theta}{(1 - \cot \theta)} = (\cos \theta + \sin \theta)$

Sol. LHS =  $\frac{\cos \theta}{1 - \tan \theta} + \frac{\sin \theta}{1 - \cot \theta}$

$$= \frac{\cos^2 \theta}{\cos \theta - \sin \theta} + \frac{\sin^2 \theta}{\sin \theta - \cos \theta} \quad 1$$

$$= \frac{\cos^2 \theta - \sin^2 \theta}{\cos \theta - \sin \theta} \quad 1$$

$$= \cos \theta + \sin \theta = \text{RHS} \quad 1$$

OR

Prove that  $(\sin \theta + \operatorname{cosec} \theta)^2 + (\cos \theta + \sec \theta)^2 = 7 + \tan^2 \theta + \cot^2 \theta$ .

Sol.  $(\sin \theta + \operatorname{cosec} \theta)^2 + (\cos \theta + \sec \theta)^2$

$$= \sin^2 \theta + \operatorname{cosec}^2 \theta + 2 + \cos^2 \theta + \sec^2 \theta + 2 \quad \frac{1}{2} + \frac{1}{2}$$

$$= \sin^2 \theta + 1 + \cot^2 \theta + 2 + \cos^2 \theta + 1 + \tan^2 \theta + 2 \quad \frac{1}{2} + \frac{1}{2}$$

$$= 7 + \tan^2 \theta + \cot^2 \theta \quad 1$$

33. In Fig. 3, XP and XQ are tangents from X to the circle with centre O. R is a point on the circle and AB is tangent at R. Prove that:

$$\mathbf{XA + AR = XB + BR}$$




Fig. 3

Sol.  $XP = XQ$  (tangents from external points) 1

$$XA + AP = XB + BQ \quad 1$$

$$XA + AR = XB + BR \quad (\text{AP} = \text{AR}, \text{BQ} = \text{BR}) \quad 1$$

34. The radii of two circles are 8 cm and 6 cm. Find the radius of the circle having its area equal to the sum of the areas of the two circles.

Sol. Let r be the radius of required circle

$$\text{Here, } \pi(8)^2 + \pi(6)^2 = \pi r^2 \quad 1$$

$$100 = r^2 \quad 1$$

$$r = 10 \text{ cm} \quad 1$$


---

### SECTION D

Question Nos. 35 to 40 carry 4 marks each.

35. In a right triangle, prove that the square of the hypotenuse is equal to sum of squares of the other two sides.

Sol. For correct given, to prove, construction and figure  $4 \times \frac{1}{2} = 2$

For correct proof 2

OR

Prove that the tangents drawn from an external point to a circle are equal in length.

Sol. For correct given, to prove, construction and figure  $4 \times \frac{1}{2} = 2$

For correct proof 2

---

36. A hemispherical depression is cut out from one face of a cubical wooden block of edge 21 cm, such that the diameter of the hemisphere is equal to edge of the cube. Determine the volume of the remaining block.

Sol. Let r be the radius of hemisphere  $\therefore r = \frac{21}{2}$  cm  $\frac{1}{2}$

$$\text{Volume of remaining block} = a^3 - \frac{2}{3}\pi r^3$$

$$= (21)^3 - \frac{2}{3}\pi \times \frac{21}{2} \times \frac{21}{2} \times \frac{21}{2} \quad 2$$

$$= 9261 \left[ 1 - \frac{\pi}{12} \right] \text{cm}^3 \quad 1$$

$$= 6853 \text{ cm}^3 \text{ (Approx.)} \quad \frac{1}{2}$$

OR

A solid metallic cylinder of diameter 12 cm and height 15 cm is melted and recast into 12 toys in the shape of a right circular cone mounted on a hemisphere of same radius. Find the radius of the hemisphere and total height of the toy, if the height of the cone is 3 times the radius.

**Sol.** Here,  $r = 6$  cm

$$\pi(6)^2 \times 15 = 12 \left[ \frac{1}{3} \pi r^2 \times 3r + \frac{2}{3} \pi r^3 \right] \quad 2$$

$$36 \times 15 = \frac{12}{3} [3r^3 + 2r^3] \quad \frac{1}{2}$$

$$9 \times 15 = 5r^3$$

$$r = 3 \text{ cm} \quad \frac{1}{2}$$

$$\text{Total height} = 12 \text{ cm} \quad 1$$

**37. The sum of first 6 terms of an A.P. is 42. The ratio of its 10th term to 30<sup>th</sup> term is 1:3. Find the first and the 13th term of the A.P.**

**Sol.** Here,  $\frac{6}{2}(2a + 5d) = 42$

$$\Rightarrow 2a + 5d = 14 \quad \dots(i) \quad 1$$

Also,

$$\frac{a + 9d}{a + 29d} = \frac{1}{3} \quad \dots(ii) \quad 1$$

$$\Rightarrow a = d \quad \frac{1}{2}$$

$$\text{Solving (i) and (ii), } 7a = 14 \quad \frac{1}{2}$$

$$\Rightarrow a = 2$$

$$d = 2 \quad \frac{1}{2}$$

$$a_{13} = a + 12d = 26 \quad \frac{1}{2}$$

OR

**Find the sum of all odd numbers between 100 and 300.**

**Sol.** Odd number between 100 to 300 are

$$101, 103 \dots 299$$

$$299 = 101 + (n - 1)2$$

$$\Rightarrow n = 100$$

$$S_n = \frac{100}{2}(101 + 299)$$

$$= 20,000$$

1

1

1


1

**38. From the top of a 7 m high building, the angle of elevation of the top of a cable tower is  $60^\circ$ , and the angle of depression of its foot is  $45^\circ$ . Find the height of the tower. Given that  $\sqrt{3} = 1.732$ .**

**Sol.**

Correct figure

1



$$\tan 45^\circ = \frac{7}{x}$$

$$\Rightarrow x = 7 \text{ m}$$

...(i)

1

$$\tan 60^\circ = \frac{h-7}{x}$$

$$x\sqrt{3} = h - 7$$

....(ii)

1

$$\text{Solving (i) and (ii) } h = 7(\sqrt{3} + 1)$$

$$= 7 \times 2.732$$

$$= 19.124 \text{ m}$$

1

**39. Find the mean of the following distribution:**

Classes:	100 – 150	150 – 200	200 – 250	250 – 300	300 – 350
Frequency:	4	5	12	2	2

**Sol.**

CI	$f_i$	$x_i$	$d_i$	$u_i$	$f_i u_i$
100-150	4	125	-100	-2	-8
150-200	5	175	-50	-1	-5
200-250	12	225	0	0	0
250-300	2	275	50	1	2
300-350	2	325	100	2	4
Total	25				-7

Correct Table

2

$$\begin{aligned} \text{Mean} &= A + \frac{\sum f_i u_i}{\sum f_i} \times h \\ &= 225 + \frac{-7}{25} \times 50 && 1 \\ &= 211 && 1 \end{aligned}$$


---

40. The sum of the reciprocals of the ages of a child 3 years ago and 5 years hence from now is  $\frac{1}{3}$ . Find his present age.

**Sol.** Let the present age = x years

$$\begin{aligned} \frac{1}{x-3} + \frac{1}{x+5} &= \frac{1}{3} && 2 \\ \Rightarrow x^2 - 4x - 21 &= 0 && 1 \\ (x-7)(x+3) &= 0 \\ \text{Hence, present age} &= 7 \text{ years} && 1 \end{aligned}$$


---